

Planar Shuffle Product, Co-Addition and the non-associative Exponential

by L. Gerritzen

Abstract

In this note we introduce the concept of a shuffle product $\sqcup$ for planar tree polynomials and give a formula to compute the planar shuffle product $S \sqcup T$ of two finite planar reduced rooted trees S, T .

It is shown that $\sqcup$ is dual to the co-addition Δ which leads to a formula for the coefficients of $\Delta(f)$.

It is also proved that $\Delta(EXP) = EXP \hat{\otimes} EXP$ where EXP is the generic planar tree exponential series, see [G]. Systems of quadratic relations for the coefficients of EXP are derived.

KEYWORDS: Planar reduced rooted trees, reduction and contraction of planar rooted trees, shuffle product, non-associative generic exponential series, co-addition, Hopf algebras

Introduction

Let $K\{x\}_\infty$ be the algebra of planar tree polynomials over a field K where the m -ary product is obtained by the K -multi-linear extension of the m -ary planar grafting of finite planar reduced rooted trees. There is a unique K -algebra homomorphism

$$\Delta : K\{x\}_\infty \rightarrow K\{x\}_\infty \otimes K\{x\}_\infty$$

mapping x onto $x \otimes 1 + 1 \otimes x$ where x denotes the tree with a single vertex. This coproduct is also called co-addition. Dually there is a shuffle product

$$\sqcup : K\{x\}_\infty \otimes K\{x\}_\infty \rightarrow K\{x\}_\infty$$

which is commutative and associative.

It can be defined by the contraction of planar trees onto sets of leaves.

For any $f \in K\{x\}_\infty$ the coefficients $(v, w(\Delta(f)))$ of $\Delta(f)$ can be computed by a formula involving the coefficients of the shuffle product $V \sqcup W$, see Prop. 4.3.

The co-addition extends continuously to a K -algebra homomorphism

$$\hat{\Delta} : K\{\{x\}\}_\infty \rightarrow K\{\{x\}\}_\infty \hat{\otimes} K\{\{x\}\}_\infty$$

of formal power series completions of $K\{x\}_\infty$ and $K\{x\}_\infty \otimes K\{x\}_\infty$.
It is shown in Section 5 that

$$\hat{\Delta}(EXP) = EXP \hat{\otimes} EXP$$

if EXP is the generic non-associative exponential series, see [G], Section 3. One obtain systems of quadratic relations for the coefficients of EXP in which the coefficients of planar shuffle products appear.

This result can be used to extend the canonical projection of [R], Chap. 3 or [L], to the non-associative setting. This application has been suggested to me by J. L. Loday.

1 Reductions of planar rooted trees

Let T be a finite planar rooted tree, see [G]. For any vertex a of T we denote by $val_T(a)$ the valence of a in T which is the number of edges of T incident with a .
Let

$$ar_T(a) := \begin{cases} val_T(a) & : \text{ if } a \text{ is the root of } T \\ val_T(a) - 1 & : \text{ otherwise} \end{cases}$$

It is called the arity of the vertex a in T .

Definition 1.1. T is called *reduced*, if $ar_T(a) \neq 1$ for all vertices a of T .

In the following construction we associate a reduced planar rooted tree $\text{Red}(T)$ to a given finite planar rooted tree T which might not be reduced.

For vertices a, b of T we denote by $[a, b]_T$ the smallest connected subgraph of T which contains a and b . It is a line and consists of the vertices and edges of the simple path between a and b .

Let

$$R^0 := \{a \in T^0 \mid \text{set of vertices of } T : ar_T(a) \neq 1\}$$

and

$$\bar{R} := \{\{a, b\} : a, b \in R^0, a \neq b, [a, b]_T^0 \cap R^0 = \{a, b\}\}.$$

Let ρ_R be the vertex of R^0 closest to the root ρ_T of T .

Proposition 1.2. $(R^0, \bar{R}, \rho_R)$ is a planar reduced rooted tree; it is called the *reduction of T* and is also denoted by $\text{Red}(T)$.

This result will follow from Proposition 1.4 below.

Remark 1.3. If the arity of the root ρ_T of a planar rooted tree T is 1, then $\text{Red}(T) = \text{Red}(T - \rho_T)$ where $T - \rho_T$ is obtained from T by deleting ρ_T and the unique edge in T incident with ρ_T .

Proposition 1.4. (i) If $T_1, \dots, T_m$ are finite planar reduced rooted trees and

$$T = \cdot_m(T_1, \dots, T_m)$$

is the grafting of $T_1, \dots, T_m$, then T is a planar reduced rooted tree if $m \geq 2$.

(ii) If $T_1, \dots, T_m$ are finite planar rooted trees and $m \geq 2$, then

$$\text{Red}(\cdot_m(T_1, \dots, T_m)) = \cdot_m(\text{Red}(T_1), \dots, \text{Red}(T_m)).$$

Proof. 1) By definition the set T^0 of vertices of T consists of the root ρ_T and the disjoint union the set T_i^0 of vertices of T_i for $1 \leq i \leq m$.

Obviously for any vertex of T_i^0 we get


$$\text{ar}_{T_i}(a) = \text{ar}_T(a)$$

while $\text{ar}_T(\rho_T) = m$.


This shows statement (i).



2) Let $T = \cdot_m(T_1, \dots, T_m)$. Then $R^0 = \{a \in T^0 : \text{ar}_T(a) \neq 1\}$ is the disjoint union $R_1^0, \dots, R_m^0$ where $R_i^0 = \{a \in T_i^0 : \text{ar}_{T_i}(a) \neq 1\}$ as the arity of the root of T is equal to $m \geq 2$. Any line $[a, b]_T$ in T for which all vertices $\neq a, \neq b$ have arity 1 in T is therefore lying in a subtree T_i of T . From this the statement (ii) follows. □

Example 1.5. The reduction T of the following planar rooted trees T_1, T_2, T_3 are all equal to $T =$



In all cases the root of the trees is the most upward vertex.





2 Contractions

Denote by **PRT** the set of isomorphism classes of finite planar reduced rooted trees.

Let $T \in \mathbf{PRT}$ and I be a subset of the set $L(T) := \{a \in T^0 : ar_T(a) = 0\}$ of leaves of T .

Then $T_I := \bigcup_{a \in I} [a, \rho_T]_T$ is a connected subgraph of T . It is a subtree of T because every connected subgraph of a tree is again a tree. The tree T_I is the smallest subtree of T containing I and ρ_T . We consider T_I as a rooted tree with root ρ_T . Then $L(T_I) = I$ because all vertices b of $[a, \rho_T]_T$ different from a have arity 1 in $[a, \rho_T]_T$.

Definition 2.1. *The contraction $T|I$ of I onto I is defined to be the reduction $Red(T_I)$ of T_I .*

Proposition 2.2. *Let $T \in \mathbf{PRT}$, $ar(T) = m \geq 2$ $T = T_1 \cdot T_2 \cdot \dots \cdot T_m$ with $T_i \in \mathbf{PRT}$.*

Let $I \subset L(T) = L(T_1) \dot{\cup} \dots \dot{\cup} L(T_m)$ and $I_j = I \cap L(T_j)$.

Then $T|I = T_1|I_1 \cdot \dots \cdot T_m|I_m$.

Proof. By standard considerations. □

3 Shuffle product

Let K be a field and $\mathbf{A} = K\{x\}_\infty$ the K -algebra with unit of planar tree polynomials over K .

A K -base of $\mathbf{A}$ is given by $\mathbf{PRT}' = \{1\} \dot{\cup} \mathbf{PRT}$ where 1 denotes the empty tree and the m -ary multiplication $\cdot_m : \mathbf{A}^m \rightarrow \mathbf{A}$ is the K -multi-linear extension of the m -ary grafting on $\mathbf{PRT}'$, see [G].

For $S, T, V \in \mathbf{PRT}'$ let $N_{S,T}(V) := \{I \subset L(V) : V|I = S, V|I^c = T\}$ where $I^c := L(V) - I$ is the complement of I in $L(V)$.

If $\deg(V) \neq \deg(S) + \deg(T)$, then $N_{S,T}(V)$ is empty.

Definition 3.1.

$$S \sqcup T := \sum_{V \in \mathbf{PRT}'} \#N_{S,T}(V) \cdot V$$

is called the planar shuffle product of S and T .

Remark 3.2. This planar shuffle product is a generalization of the usual shuffle product as defined in [R], section (1.4).

Proposition 3.3. *There is a unique K -bilinear map*

$$\sqcup : K\{x\}_\infty \times K\{x\}_\infty \rightarrow K\{x\}_\infty$$

such that $\sqcup(S, T)$ is the shuffle product of S and T for any $S, T \in \mathbf{PRT}'$.

It is called the shuffle product on $K\{x\}_\infty$.

Moreover $\sqcup$ is commutative and associative.

Proof. 1) The existence and uniqueness of the K -bilinear map follows immediately from the fact that $\mathbf{PRT}'$ is a K -base of $K\{x\}_\infty$.

2) The commutativity of $\sqcup$ follows because $N_{S,T}(V) = N_{T,S}(V)$. A bijection between $N_{S,T}(V)$ and $N_{T,S}(V)$ is given by $I \rightarrow I^c$.

3) Let now $R, S, T \in \mathbf{PRT}'$ and $N_{R,S,T}(V) := \{(I_1, I_2, I_3) : I_j \subseteq L(V) \text{ } I_1, I_2, I_3 \text{ is a disjoint union of } L(V), V|_{I_1} = R, V|_{I_2} = S, V|_{I_3} = T\}$. It is an easy exercise to show that

$$R \sqcup S) \sqcup T = \sum_{W \in \mathbf{PRT}'} \#N_{R,S,T}(W) \cdot W.$$

The right hand side is also equal to $R \sqcup (S \sqcup T)$. □

For $k \in \mathbb{N}_{\geq 1}$ denote by $\underline{k}$ the set $\{1, 2, \dots, k\}$ of integers between 1 and k .

For any non-empty finite subset α of $\mathbb{N}_{\geq 1}$ denote by $\alpha[i]$ the i -th element of α relative to the natural order on α for $i \in \underline{k}$, $k := \#\alpha$. Thus the map

$$i \mapsto \alpha[i]$$

is a bijection between $\underline{k}$ and α and $\alpha[i] < \alpha[i+1]$ for $1 \leq i < k$.

We represent α by the word $\alpha[1] \cdot \alpha[2] \cdot \dots \cdot \alpha[k]$.

For any $k \in \mathbb{N}_{\geq 2}$ denote by $\Gamma_k := \{(\alpha, \beta) : \alpha \subseteq \underline{k}, \beta \subseteq \underline{k}, \alpha \neq \emptyset \neq \beta, \alpha \cup \beta = \underline{k}\}$ the set of pairs (α, β) of non-empty subsets of $\underline{k}$ whose union is the whole set $\underline{k}$. For any $m, n \in \mathbb{N}_{\geq 1}$ denote $\Gamma_k(m, n) := \{(\alpha, \beta) \in \Gamma_k : \sharp\alpha = m, \sharp\beta = n\}$. Obviously the map $(\alpha, \beta) \mapsto (\beta, \alpha)$ is a bijection between $\Gamma_k(m, n)$ and $\Gamma_k(n, m)$.

Lemma 3.4. *i) $\Gamma_k(m, n) \neq \emptyset$ if and only if $m \leq k, n \leq k, k \leq m + n$.*

ii) If $m \leq k, n \leq k, k \leq m + n$, then $\sharp\Gamma_k(m, n) = \frac{k!}{(k-n)!(k-m)!(m+n-k)!}$

Proof. 1) (i) is immediate, as for $(\alpha, \beta) \in \Gamma_k(m, n)$ one has $\alpha \subseteq \underline{k}, \beta \subseteq \underline{k}, \alpha \cup \beta = \underline{k}$. from which $\sharp\alpha \leq k, \sharp\beta \leq k, \sharp\alpha + \sharp\beta \geq \sharp\underline{k} = k$ follows.

2) Assume the assumption of (ii). There are $\binom{k}{n}$ subsets α of $\underline{k}$ with n elements. There are $\binom{n}{m+n-k}$ subsets γ of α with $m + n - k$ elements. Let $\beta := (\underline{k} - \alpha) \cup \gamma$. Any pair of $\Gamma_k(m, n)$ is obtained by this procedure. Thus $\sharp\Gamma_k(m, n) = \binom{k}{n} \cdot \binom{n}{m+n-k}$.

□

Example 3.5. $\Gamma_2(1, 1) = \{(1, 2), (2, 1)\}$

$\Gamma_2(1, 2) = \{(1, 12), (2, 12)\}$

$\Gamma_2(2, 1) = \{(12, 1), (12, 2)\}$

$\Gamma_2(2, 2) = \{(12, 12)\}$

$\Gamma_2(m, n) = \emptyset$ if m or n is ≥ 3

$\Gamma_k(1, n) \neq \emptyset$ if and only if $n \leq k \leq n + 1$.

$\Gamma_n(1, n) = \{(i, \underline{n}) : i \in \underline{n}\}$

$\Gamma_{n+1}(1, n) = \{(i, \underline{n+1} - \{i\}) : i \in \underline{n}\}$.

Especially

$\Gamma_3(1, 3) = \{(1, \underline{3}), (2, \underline{3}), (3, \underline{3})\}$

$\Gamma_4(1, 4) = \{(1, 234), (2, 134), (3, 124), (4, 123)\}$.

Let $\bar{\Gamma}_k(m, n) := \Gamma_k(1, 1) \cup \Gamma_k(m, 1) \cup \Gamma_k(1, n) \cup \Gamma_k(m, n)$.

It is the set of all pairs (α, β) of subsets of $\underline{k}$ such that $(\sharp\alpha$ is 1 or $m, \sharp\beta$ is 1 or n and $\alpha \cup \beta = \underline{k}$.

Let $S, T \in \mathbf{PRT}, m = ar(S), n = ar(T), S = S_1 \cdot S_2 \cdot \dots \cdot S_m, T = T_1 \cdot T_2 \cdot \dots \cdot T_n$ with $S_i, T_j \in \mathbf{PRT}$.

For any $(\alpha, \beta) \in \bar{\Gamma}_k(m, n)$ we are going to define a polynomial $R_{(\alpha, \beta)} \in K\{x\}_\infty$. One has to consider several cases:

Case 1:

$\sharp\alpha = \sharp\beta = 1$. Then $(\alpha, \beta) \in \Gamma_2(1, 1)$ and $R_{(\alpha, \beta)} := S \cdot T$ if $\alpha = 1, \beta = 2$ and $R_{\alpha, \beta} := T \cdot S$ if $\alpha = 2, \beta = 1$.

Case 2:

$$\sharp\alpha = 1, \sharp\beta = m > 1, \alpha = \{\nu\}.$$

Then $(\alpha, \beta) \in \Gamma_k(1, n)$ and k is either n or $n + 1$.

If $k = n + 1$, then $\beta = \underline{n+1} - \{\nu\}$ and

$$R_{(\alpha, \beta)} := T_1 \cdot T_2 \cdot \dots \cdot T_{\nu-1} \cdot S \cdot T_\nu \cdot T_{\nu+1} \cdot T_n$$

which is a tree of arity $n + 1$.

If $k = n$, then $\beta = \underline{n}$ and $R_{(\alpha, \beta)} := T_1 \cdot T_2 \cdot \dots \cdot T_{\nu-1} \cdot (T_\nu \sqcup S) \cdot T_{\nu+1} \cdot \dots \cdot T_n$
which is a polynomial of arity n .

Case t :

$\sharp\beta = 1, \beta = \{\mu\}, \sharp\alpha = n > 1$. Then $(\alpha, \beta) \in \Gamma_k(m, 1)$ and $k = m$ or $m + 1$.

If $k = m + 1$, then $\alpha = \underline{m+1} - \{\mu\}$ and

$R_{(\alpha, \beta)} := S_1 \cdot S_2 \cdot \dots \cdot S_{\mu-1} \cdot T \cdot S_\mu \cdot S_{\mu+1} \cdot \dots \cdot S_m$ which is a tree of arity $m + 1$.

If $k = m$, then $\alpha = \underline{m}$ and $R_{(\alpha, \beta)} := S_1 \cdot S_2 \cdot \dots \cdot S_{\mu-1} \cdot (S_\mu \sqcup T) \cdot S_{\mu+1} \cdot \dots \cdot S_m$
which is a polynomial of arity m .

Case 3:

$\sharp\alpha = m > 1, \sharp\beta = n > 1$. Then $R_{(\alpha, \beta)} := P_1 \cdot P_2 \cdot \dots \cdot P_k$

where $P_j = S_i$ if $j \in \alpha, j \notin \beta$ and $\alpha[i] = j$

where $P_j = T_i$ if $j \in \beta, j \notin \alpha$ and $\beta[i] = j$

and where $P_j = S_i \sqcup T_l$ if $j \in \alpha \cap \beta$ and $\alpha[i] = j, \beta[l] = j$.

Proposition 3.6.

$$S \sqcup T = \sum_{k=2}^{n+m} \sum_{(\alpha, \beta) \in \bar{\Gamma}_k(m, n)} R_{(\alpha, \beta)}.$$

Proof. 1) Let $N_{S, T} = \cup_{V \in \mathbf{PRT}} N_{S, T}(V)$.

Its elements can be seen as pairs (V, I) with $V \in \mathbf{PRT}$, $I \subset L(V)$ and $I \in N_{S, T}(V)$. Obviously $S \sqcup T = \sum V$ where the summation is extended over $N_{S, T}$.

- 2) Let $(V, I) \in N_{S,T}$ and V be of arity k . Then $V = V_1 \cdot V_2 \cdot \dots \cdot V_k$ with $V_i \in \mathbf{PRT}$ and $L(V) = L(V_1) \dot{\cup} \dots \dot{\cup} L(V_k)$.

The type (α, β) of (V, I) is defined as follows:

$$\alpha := \{i \in \underline{k} : I \cap L_i(V) \neq \emptyset\}$$

$$\beta := \{i \in \underline{k} : I^c \cap L_i(V) \neq \emptyset\}$$

where I^c is the complement of I in $L(V)$.

It is easy to see that $(\alpha, \beta) \in \bar{\Gamma}_k(m, n)$. If $\alpha \cap \beta = \emptyset$ it is obvious that $V = R_{(\alpha, \beta)}$ and $I = \cup_{i \in \alpha} L_i(V)$.

- 3) If $\alpha \cap \beta \neq \emptyset$ then the situation is more complicated to describe.

What happens can already be seen if $m = n = k$ and $\alpha = \beta = \underline{k}$.

Then $R_{(\alpha, \beta)} = (S_1 \sqcup T_1) \dots / S_n \sqcup T_n = \sum V_1 \cdot V_2 \cdot \dots \cdot V_n$ where the summation is extended over the set of n -tuples $((V_1, I_1), \dots, (V_n, I_n))$ with $(V_i, I_i) \in N_{S_i, T_i}$ where $S = S_1 \cdot S_2 \dots S_n, T = T_1 \cdot T_2 \cdot \dots \cdot T_n$.

This is obviously equal to the sum $\sum_{(V, I) \in N_{S, T}} V$, where $N'_{S, T}$ is the subset of $N_{S, T}$ consisting of all (V, I) for which the type is equal to $(\underline{n}, \underline{n})$.

- 4) The general case follows by a combination of the arguments in 2) and 3). $\square$

Example 3.7. $x \sqcup x = 2x^2$.

If $S = x$, and $n \geq 2$, then

$$x \sqcup T = x \cdot T + t \cdot x + \sum_{i=1}^n T_1 \cdot \dots \cdot T_{i-1} \cdot x \cdot T_i \cdot T_{i+1} \cdot \dots \cdot T_n + \sum_{i=1}^n T_1 \cdot \dots \cdot T_{i-1} \cdot (x \sqcup T_i) \cdot T_{i+1} \cdot \dots \cdot T_n.$$

It follows from the Proposition because $\bar{\Gamma}_k(1, n) = \Gamma_k(1, 1) \cup \Gamma_k(1, n)$ and from the explicit description of $\Gamma_k(1, n)$ above.

4 Co-Addition

Let $\mathbf{A} = K\{x\}_\infty$ and $B = \mathbf{A} \otimes_K \mathbf{A}$.

Proposition 4.1. *For any $m \in \mathbb{N}_{\geq 2}$ there is a unique K -multi-linear map*

$$\cdot_m : \mathbf{B}^m \rightarrow \mathbf{B}$$

such that $\cdot_m(f_1 \otimes g_1, \dots, f_m \otimes g_m) = (f_1 \cdot f_2 \cdot \dots \cdot f_m) \otimes (g_1 \cdot g_2 \cdot \dots \cdot g_m)$ for all $f_i, g_i \in \mathbf{A}$.

$\mathbf{B}$ together with this string $(\cdot_m)_{m \geq 2}$ of operations is considered as an algebra.

Proof. As in the classical case of one binary multiplication. $\square$

Proposition 4.2. *There is a unique unital K -algebra homomorphism*

$$\Delta : \mathbf{A} \rightarrow \mathbf{A} \otimes \mathbf{A}$$

such that $\Delta(x) = x \otimes 1 + 1 \otimes x$.

Δ is called co-addition or coproduct induced by addition.

Proof. By standard methods. $\square$

For $T \in \mathbf{PRT}'$ let

$$c_T : \mathbf{A} \rightarrow K$$

be the K -linear map with

$$c_T(S) = \delta_{ST} = \begin{cases} 1 & : S = T \\ 0 & : S \neq T \end{cases}$$

for any $S \in \mathbf{PRT}'$.

It is called the coefficient map relative to T .

For $V, W \in \mathbf{PRT}'$ let

$$c_{V,W} : \mathbf{A} \otimes \mathbf{A} \rightarrow K$$

be the K -linear map such that

$$c_{V,W}(S \otimes T) = \begin{cases} 1 & : (V, W) = (S, T) \\ 0 & : \text{otherwise} \end{cases}$$

for $S, T \in \mathbf{PRT}'$. It is called the coefficient map relative to $V \otimes W$.

The following formula for planar polynomials generalizes the classical formula, see [R].

Proposition 4.3. *For any $f \in K\{x\}_\infty$ and $V, W \in \mathbf{PRT}'$ one gets*

$$c_{V,W}(\Delta(f)) = \sum_{T \in \mathbf{PRT}'} c_T(f) \cdot c_T(V \sqcup W).$$

Proof. 1) For any $T \in \mathbf{PRT}'$ we obtain $\Delta(T) = \sum_{I \in L(T)} (T|I) \otimes (T|I^c)$ as can be seen by induction on $\deg(T)$, see also [GH]. It follows that $c_{V,W}(\Delta(T)) = c_T(V \sqcup W)$. Then the formula is proved if f is a monomial.

2) Both sides of the formula are K -linear and thus they are equal because of step 1). $\square$

5 Co-Addition of EXP

Let $\hat{\mathbf{A}}$ be the x -adic completion of $\mathbf{A} = K\{x\}_\infty$ and $\hat{\mathbf{B}}$ be the $(x \otimes 1, 1 \otimes x)$ -adic completion of $\mathbf{B}$. Then $\Delta : \mathbf{A} \rightarrow \mathbf{A} \otimes \mathbf{A}$ extends to a continuous coproduct

$$\hat{\Delta} : \hat{\mathbf{A}} \rightarrow \hat{\mathbf{B}} = \hat{\mathbf{A}} \hat{\otimes} \hat{\mathbf{A}}.$$

Proposition 5.1. *Let EXP be the generic exponential in $K\{\{x\}\}_\infty$, see [G], $K = \mathbb{Q}(q)$ field of rational functions in q over $\mathbb{Q}$. Then*

$$\hat{\Delta}(EXP) = EXP \hat{\otimes} EXP$$

Proof. 1) Let $k \in \mathbb{N}_{\geq 2}$ and $exp_k(x)$ be the k -ary exponential series in $\mathbb{Q}\{\{x\}\}_\infty$. Let $f = \hat{\Delta}(exp_k(x))$.

We want to show that $f = exp_k(x) \hat{\otimes} exp_k(x)$.

2) For $k \in \hat{\mathbf{B}}$ let $ord_{\mathbf{B}}(k) := \min\{deg(V) + deg(W) : c_{V,W}(h) \neq 0, V, W \in \mathbf{PRT}'\}$. Obviously $ord_{b_f \hat{\mathbf{B}}} \Delta(g) = ord_{\hat{\mathbf{A}}}(g)$ for $g \in \hat{\mathbf{A}}$ and thus

$$ord_{\hat{\mathbf{B}}}(f - (1 \otimes 1 + x \otimes 1 + 1 \otimes x)) \geq 2$$

3) There is a continuous K -automorphism

$$\varphi_k : \hat{\mathbf{B}} \rightarrow \hat{\mathbf{B}}$$

such that

$$\begin{aligned} \varphi_k(x \hat{\otimes} 1) &= k(x \hat{\otimes} 1) \\ \varphi_k(1 \hat{\otimes} x) &= k(1 \hat{\otimes} x). \end{aligned}$$

Fix k and let $f = \Delta(EXP^k) = \Delta(EXP(kx)) = \varphi_k(f)$.

Denote by f_r the homogeneous part of f degree r .

Then

$$\sum_{\substack{V, W \in \mathbf{PRT}' \\ deg V + deg W = r}} c_{V,W}(f) \cdot V \otimes W.$$

As $\varphi_k(f_r) = k^r f_r$ we get from the functional equation for f that

$$(f^k)_r = k^r f_r$$

if $(f^k)_r$ is the homogeneous part of f^k of degree r which is

$$\sum_{\deg V + \deg W = r} c_{V,W}(f^k) \cdot V \otimes W.$$

As in [G] the Prop. (3.1) one can show that f must be equal to $\exp_k \hat{\otimes} \exp_k$.

- 4) Let R be the subring of all rational function f in $\mathbb{Q}(q)$ whose poles are roots of unity $\neq 1$.

Let $\pi_k : R \rightarrow \mathbb{Q}$ be the $\mathbb{Q}$ -algebra homomorphism with $\pi_k(q) = k$ for $k \in \mathbb{N}_{\geq 2}$.

Now π_k induces algebra homomorphisms

$$\hat{\pi}_k : R\{\{x\}\}_\infty \rightarrow \mathbb{Q}\{\{x\}\},$$

$$\hat{\pi}_k^2 : R\{\{x\}\}_\infty \hat{\otimes} R\{\{x\}\}_\infty \rightarrow \mathbb{Q}\{\{x\}\}_\infty \hat{\otimes} \mathbb{Q}\{\{x\}\}_\infty$$

and

$$\hat{\pi}_k^2(\Delta(k)) = \Delta(\hat{\pi}_k(k))$$

for $k \in R\{\{x\}\}_\infty$.

As $\hat{\pi}_k(EXP) = \exp_k$ and $\hat{\pi}^{(2)}(\Delta(EXP)) = \exp_k \hat{\otimes} \exp_k$ we get

$$\Delta(EXP) = EXP \hat{\otimes} EXP.$$

□

Let $a(T) = c_T(EXP)$ be the coefficient of EXP relative to $T \in \mathbf{PRT}'$. Recall that $a(1) = a(x) = 1$ and the recursive relation

$$a(T) = \frac{\binom{q}{m}}{q^n - q} a(T_1) \dots a(T_m)$$

if $m = ar(T)$, $T = T_1 \cdot \dots \cdot T_m$ and $n = \deg(T)$, see [G].

In the following Corollary one obtains a quadratic relation for the system $(a_T)_{T \in \mathbf{PRT}'}$.

Corollary 5.2. $a(V) \cdot a(W) = \sum_T a_T c_T(V \sqcup W).$

Example 5.3. Let $m \geq 3$ and $S = x \sqcup x^m$. One can show that $S = (m+1)x^{m+1} + 2(x^2 \cdot x \cdot \dots \cdot x + x \cdot x^2 \cdot x \cdot \dots \cdot x + \dots + x \cdot x \cdot \dots \cdot x \cdot x^2) + x \cdot x^m + x^m \cdot x$ by applying Example (3.6).

From the recursive formula for the coefficients of EXP one gets

$$a(x^m) = \frac{\binom{q}{m}}{q^m - q}$$

$$a(x^2 \cdot x \cdot \dots \cdot x) = \frac{\binom{q}{m}}{q^{m+1} - q} \cdot \frac{1}{2} = a(x \cdot \dots \cdot x \cdot x^2 \cdot x \cdot \dots \cdot x)$$

if the factor x occurs m times, as $a(x^2) = \frac{1}{2}$.

$$a(x^m \cdot x) = a(x \cdot x^m) = \frac{\binom{q}{2}}{q^{m+1} - q} \cdot \frac{\binom{q}{m}}{q^m - q}.$$

The right hand side in the formula of Corollary (5.2) is

$$b = \sum_{T \in \mathbf{PRT}'} c_T(S) \cdot a(T).$$

We are substituting the values above for $a(T)$ and obtain

$$b = (m+1) \frac{\binom{q}{m+1}}{q^{m+1} - q} + 2m \frac{\binom{q}{m}}{q^{m+1} - q} \cdot 1/2 + 2 \frac{\binom{q}{2}}{q^{m+1} - q} \frac{\binom{q}{m}}{q^m - q}$$

as $\deg((x^{m+1})) = m+1$, $\deg(x^2 \cdot x \cdot \dots \cdot x) = m+1$ if the factor x occurs m times and $\deg(x \cdot x^m) = \deg(x^m \cdot x) = m+1$.

Thus

$$b = \frac{\binom{q}{m}}{q^m - q} ((m+1) \frac{\binom{q-m}{m+1} \cdot (q^m - q)}{(q^{m+1} - q)} + m \cdot \frac{q^m - q}{q^{m+1} - q} + \frac{q(q-1)}{q^{m+1} - q}) = \frac{\binom{q}{m}}{q^m - q}$$

because

$$(q-m)(q^m - q) + m(q^m - q) + q(q-1) = q^{m+1} - q$$

and $b = a(x) \cdot a(x^m) = a(x^m) = \frac{\binom{q}{m}}{q^m - q}$ which is in accordance with Corollary (5.2).

References

- [DG] Drensky, V. - Gerritzen, L.: Non-associative Exponential and Logarithm, Nonassociative exponential and logarithm, Journal of Algebra 272 (2004) 331 - 320
- [G] Gerritzen, L.: Planar rooted trees and non-associative exponential series, Advances in Applied Mathematics 33 (2004) 342 - 365
- [GH] Gerritzen, L. - Holtkamp, R.: Hopf Co-Addition for Free Magma Algebras and the non-associative Hausdorff series, Journal of Algebra 265(2003), 264-284
- [L] Loday, J.-L. : Série de Hausdorff, idempotents Eulériens et algèbres de Hopf, Expo. Math 12(1994) 165-178
- [R] Reutenauer, C.: Free Lie Algebras, Oxford Univ. Press, 1993